


Transmission Lines, Greater Sage-Grouse, Ravens, and Raptors in Southwestern Idaho







OBJECTIVES

- Review information on the response of raptors and ravens to new transmission lines
- Discuss information on nest predation of greater-sage grouse by ravens
- Predict the effects of proposed transmission lines in southwestern Idaho on raptors, ravens, and sage-grouse




Snake River Birds of Prey







Number of Raptors and Raven Pairs Nesting on the PP&L 500-kV line




Steenhof, K., M.N. Kochert and J.A. Roppe. 1993. Nesting by raptors and common ravens on electrical transmission line towers. Journal of Wildlife Management 57: 271-281.











Engel, K.A., L.S. Young, K. Steenhof, J.A. Roppe and M.N. Kochert. 1992. Communal roosting of common ravens in southwestern Idaho. Wilson Bulletin 104: 105-121.




Daily Movements of Common Ravens

Maximum Distance from Roost Mean = 7 km Maximum = 65 km

Engel, K.A. and L.S. Young. 1992. Movements and habitat use by common ravens from roost sites in southwestern Idaho. Journal of Wildlife Management 56: 596-602.

IMPLICATIONS?







- Nest failure is an important factor in sage-grouse declines
- Nest predation is the primary cause of nest failure
- Ravens are the most common nest predator in NE Nevada

Peter S. Coates, John W. Connelly, and David J. Delehanty. 2008. Predators of Greater Sage-Grouse nests identified by video monitoring. *J. Field Ornithol.* 79(4):421–428

GATEWAY WEST Transmission Line Project



Office / Idal




Snake River Birds of Prey

National Conservation Area


Resource Management Plan and Record of Decision


September 2008








GATEWAY WEST Transmission Line Project






























SUMMARY



- Raptors and ravens will be attracted to new transmission lines.
- Colonization will occur mainly near existing breeding populations.
- Buffers around leks and brood-rearing areas need to be large to protect grouse from avian predation.
- Transmission lines can be compatible with raptors.



SITE-SPECIFIC CONCLUSIONS

- One of two routes proposed in the NCA would be compatible with raptors and NCA goals
- It would avert an alternative that would have adverse effects on Greater Sage-Grouse.
- The other route has issues and conflicts that need to be resolved.

